

Cupaniopsis anacardioides carrotwood



Appearance: Slender evergreen tree, usually single-trunked, to 10 m (33 ft) tall, with dark gray outer bark and often orange inner bark.

Leaves: Alternate, once compound (usually even-pinnate), with petioles swollen at the base; 4-12 leaflets, stalked, oblong, leathery, shiny yellowish green, to 20 cm (8 in) long and 7.5 cm (3 in) wide, with margins entire and tips rounded or slightly indented.


Flowers: Numerous, white to greenish yellow, up to 0.8 cm (0.4 in) wide, in branched clusters to 35 cm (14 in) long at leaf axils; 5 petals; 6-8 stamens.

Fruit: A short-stalked, woody capsule, to 2.2 cm (0.9 in) across, with 3 distinctly ridged segments; yellow orange when ripe, drying to brown and splitting open to expose 3 shiny oval black seeds covered by a yellow-red crust (aril).

Ecological threat:

Invades spoil islands, beach dunes, marshes, tropical hammocks, pinelands, mangrove and cypress swamps, scrub habitats, and coastal strands; greatly altering understory habitat. FLEPPC Category I

Distribution: NW, NE, C, SW, SE



Field Notations

CUPAANAC/CUAN4

<http://plants.ifas.ufl.edu>